Journal of Immunological Methods, 168 (1994) 285–286
Elsevier Science B.V.

Contents to volume 168 (1994)

Number 1 – 12 January 1994

Research reports

A new human cell line for the molecular dissection of the requirement for interleukin-1 in the production of interleukin-2 by immature T cells
U.R. Kees, S.E. Peroni, P.R. Ranford and J. Ford (Australia) 1

Quantification of recombinant human interleukin-1α by a specific two cell immunobioassay
R.W. Nadeau, C.M. Ostrowski, G. Ni-Wu and D.J. Liberato (USA) 9

Development of an ELISA for measuring the activity of tetanus toxoid in vaccines and comparison with the toxin neutralization test in mice
M.A. Manghi, M.F. Pasetti, M.L. Brero, S. Deluchi, G. Di Paola, V. Mathet and P.V. Eriksson (Argentina) 17

Restoration of a normal level of immunoglobulin production in a hybridoma cell line following modification of the chromosomal immunoglobulin μ gene by gene replacement
M.D. Baker, H.A. Karn and L.R. Read (Canada) 25

The effect of α_2 macroglobulin in commercial cytokine assays
K. James, I. Milne, A. Cunningham and S.-F. Elliott (UK) 33

A new, simple, bioassay for human IFN-γ
W. Däubener, N. Wanagat, K. Pilz, S. Seghrouchni, H.G. Fischer and U. Hadding (Germany) 39

A sensitive and specific bioassay for the detection of human interleukin-10
J.F. Schlaak, E. Schmitt, C. Hüls, K.-H. Meyer zum Büschenfelde and B. Fleischer (Germany) 49

Detection of anti-helminth antibody by microenzyme-linked immunosorbent assay using recombinant antigen and anti-β-galactosidase monoclonal antibody
K. Sugane and S.-H. Sun (Japan) 55

A novel and sensitive method for the detection of secreted cell products using time-resolved fluorescence
C. Ruedl, G. Wick and H. Wolf (Austria) 61

Monoclonal antibodies as probes to detect conformational changes in the rat cysteine proteinase inhibitor cystatin A
A. Takeda, A. Iwasawa, Y. Nakamura, K. Omata and K. Nakaya (Japan) 69

Production of lymphotoxin (LTα) and a soluble dimeric form of its receptor using the baculovirus expression system
P.D. Crowe, T.L. VanArsdale, B.N. Walter, K.M. Dahms and C.F. Ware (USA) 79

Affinity ranking of influenza neuraminidase mutants with monoclonal antibodies using an optical biosensor. Comparison with ELISA and slot blot assays
L.C. Gruen, J.L. McKimm-Breschkin, J.B. Caldwell and E.C. Nice (Australia) 91

Influence of solid-phase antigen in competition enzyme-linked immunosorbent assays (ELISAs) on calculated antigen-antibody dissociation constants
S.J. Seligman (USA) 101

An ultrasensitive chemiluminoenzyme immunoassay for the quantification of human tissue kininogens: application to synovial membrane and cartilage
F. Legris, J. Martel-Pelletier, J.-P. Pelletier, R. Colman and A. Adam (Canada, USA) 111

Induction of plasmacytomas that secrete monoclonal anti-peptide antibodies by retroviral transformation
E.M. Weissinger, D.W. Henderson, H. Mischak, J.A. Goodnight and J.F. Mushinski (USA) 123

Production of large amounts of recombinant interleukins by cDNA transfected mouse myeloma cells cultured in dialysis tubing
E. Sjögren-Jansson, C. Gustafsson, S. Jeansson, U. Karlsson and N. Lycke (Sweden) 131

A high-capacity manifold support for the detection of specific IgE antibodies in allergic individuals
M. Kwiatkowski, J. Parik and U. Landegren (Sweden) 137

Book reviews 145

Notes to authors 147

Number 2 – 10 February 1994

Minireview

The genetic engineering of monoclonal antibodies
R.J. Owens and R.J. Young (UK) 149

Research reports

Detection of allergen- and mitogen-induced human cytokine transcripts using a competitive polymerase chain reaction
S.-K. Huang, D.M. Essayan, G. Krishnaswamy, M. Yi, M. Kumai, S.-N. Su, H.-Q. Xiao, L.M. Lichtenstein and M.C. Liu (USA) 167

A simplified rapid method for restriction fragment length polymorphism analysis after bone marrow transplantation
T. Schreiner, M. Wiesneth, B. Maccari, B. Sawodny, H. Prochnow-Calzia and B. Kubanek (Germany) 183

Ammonium chloride exposure inhibits cytokine-mediated eosinophil survival
M. Ide, D. Weiler, H. Kita and G.J. Gleich (USA) 187

A simple method for the immunocytochemical processing of large numbers of floating sections and its application in screening for monoclonal antibodies
L.R. Berghman, L. Grauwels, L. Vanhamme and F. Vandesande (Belgium) 197

Flow cytometric measurement of non-specific esterase activity
F. Cook and L.-J. Eales (UK) 203

Non-radioactive labelling and immunoprecipitation analysis of leukocyte surface proteins using different methods of protein biotinylation
T. Kähne and S. Ansorge (Germany) 209

Measuring CDR3 length variability in individuals during ontogeny
S. Desravines and E. Hsu (USA) 219

Effect of antibody density on the displacement kinetics of a flow immunoassay
S.Y. Rabbany, A.W. Kusterbeck, R. Bredehorst and F.S. Ligler (USA, Germany) 227

Riboprobe expression cassettes for measuring IGF-I, β-actin and glyceraldehyde 3-phosphate dehydrogenase transcripts
A. Biragyn, S. Arkins and K.W. Kelley (USA) 235

Use of Merocyanine 540 and Hoechst 33258 for the selective killing of contaminating mycoplasmas in cell cultures
M. Tarshis, A. Katzenel and S. Rottem (Israel) 245

An improved MTT assay using the electron-coupling agent menadione
H. Garn, H. Krause, V. Enzmann and K. Dröβler (Germany) 253

Typing of the HLA-DRB3 gene by temperature gradient gel electrophoresis. Prediction of the resolution of four allelic fragments by computational simulation of DNA melting
M. Chen, W. März, K.H. Usadel, E.H. Scheuermann and B.O. Boehm (Germany) 257

Simultaneous measurement of natural killer cell cytotoxicity against each of three different target cell lines
K. Blomberg (Finland) 267

Identification of a higher molecular weight protein that shows apparent cross-reactivity with anti-p21^{ras} monoclonal antibodies on Western blots
Z. Xiang, X. Wang, J. Gao, K.J. Morrow, Jr. and R.A. Nakashima (USA) 275

Letter to the editors

Ability of 'OptiLyse' lysing/fixing reagents to inactivate HIV-infected H9 cells in whole blood
J.K.A. Nicholson and S.W. Browning (USA) 283

Contents to volume 168 (1994) 285

Notes to authors 287